

This open-file release is a progress report that provides to the public the results of the first year of mapping on a three-year project. The map is incomplete and inconsistencies, errors, and omissions have not been resolved. This map may not conform to UGS policy and editorial standards and it may be premature for an individual or group to take action based on the contents.

The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either express or implied, of the U.S. Government.

Progress Report Geologic Map of the Dutch John Quadrangle (east part)
Utah-Colorado-Wyoming
Year 1 of 3

Compiled by Douglas A. Sprinkel
2000

Utah Geological Survey
a division of
Utah Department of Natural Resources
in cooperation with
U.S. Geological Survey
STATEMAP Agreement No. 99HQAG0138

30 X 60 MINUTE SERIES (TOPOGRAPHIC)



SCALE 1:100 000

CENTIMETER ON THE MAP REPRESENTS 1 KILOMETER ON THE GROUND
CONTOUR INTERVAL 50 METERS

DUTCH JOHN, UTAH-COLO.-WYO.
N4030-W10900/30 X 60

Tcd	Chinle, Moenkopi, and Dinwoody Formations undivided (Upper and Lower Triassic) - Chinle, Moenkopi, and Dinwoody Formations are combined as a single map unit on the north flank of the Uinta Mountains; On previous maps, Chinle beds have been called the Ankaresh Formation and the Moenkopi beds have been called the Woodside Shale on the north flank of the Uinta Mountains. Chinle Formation - See description below; 91-116 m Moenkopi Formation - See description below; 221 m thick Dinwoody Formation - Light-gray, greenish-gray, light-brown, and brown, thin-bedded ripple-marked shale, siltstone, and sandstone with minor amounts of limestone. Mostly a soft, slope-forming unit mapped along the north flank of the Uinta Mountains; 110-162 m thick.
Tcr	Chinle Formation (Upper Triassic) - Purplish-red, purple, light-gray, greenish-gray, light-green, rippled-bedded siltstone, sandstone, claystone, shale, and conglomerate; generally forms slopes; base is resistant conglomerate unit named the Gartra Member; 50-90 m thick.
Tm	Moenkopi Formation (Lower Triassic) - Medium- to dark-red, reddish-brown, green, and gray ripple-marked siltstone, fine-grained sandstone, and shale with gypsum and limestone beds; mostly soft, slope-forming unit; 220-240 m thick.
Ppc	Park City and Phosphoria Formations (Lower Permian) - Franson Member of Park City Formation - Gray, thick- to thin-bedded cherty limestone and dolomite interbedded with brownish-gray sandstone and red to ochre shale; generally resistant and form ledges and cliffs. Meade Peak Phosphatic Shale Member of the Phosphoria Formation - Slope-forming dark-gray phosphatic shale with interbeds of sandstone and limestone. Grandeur Member of Park City Formation - Light-gray to light-brownish-gray sandstone, dolomite, and limestone; generally resistant and form ledges and cliffs. Combined thickness of Park City and Phosphoria Formations is 73-122 m.
PIpw	Weber Sandstone (Lower Permian to Middle Pennsylvanian) - Light-gray to yellowish-gray, very thick bedded sandstone with interbeds of limestone in the lower part; highly cross-bedded sandstone in the upper part; forms steep cliffs and ridges; 472 m thick.
IPMu	Pennsylvanian and Mississippian rocks undivided - Small fault blocks of carbonate rocks along the Uinta fault zone.
IPm	Morgan Formation (Middle Pennsylvanian) - Light- to medium-red, yellow, and gray shale and siltstone, light- to medium-gray fossiliferous and red cherty limestone, and fine-grained, locally cross-bedded sandstone and red shale; 11-37 m thick.
IPrv	Round Valley Limestone (Lower Pennsylvanian) - Light-gray to light-blue-gray, thin- to very thick bedded limestone interbedded with soft red shale; limestone is fossiliferous and cherty; chert is blue gray and yellowish gray; but red to pink Jasperoid chert is common in the region; forms ledges and cliffs; 80-127 m thick
Mdh	Doughnut Shale and Humburg Formation Doughnut Shale (Upper Mississippian) - Dark-gray shale with some red shale near base with beds of coarse sandstone, limestone and coal; shale is slope forming and clayey; 91 m thick. Humburg Formation (Upper Mississippian) - Light-gray to red, fine-grained to very fine grained, soft to resistant sandstone interbedded with light-gray limestone and red to black shale; sandstone is locally cross-bedded and hematitic near the top of the formation; 75-90 m thick.
Mm	Madison Limestone (Upper and Lower Mississippian) - Mostly dark-gray, medium to coarse crystalline, cherty limestone; chert is typically light gray; commonly contains solution cavities; 130-300 m thick.
CI	Lodore Formation (Upper Cambrian) - Light-brown to greenish-gray sandstone underlain by pink to tan to pale-greenish-gray glauconitic shale interbedded with tan to pale-green sandstone; base is variegated (pink, gray, and pale-green) coarse- to medium-grained cross-bedded sandstone; locally pebbly; upper part forms ledges, middle part forms slopes and ledges, and lower part forms cliffs; 180 m thick.
Yu	Uinta Mountain Group (Upper and Middle Proterozoic) - Dark- to light-red, medium- to coarse-grained, massive to cross-bedded siliceous sandstone (metagartzite); contains considerable red, green, and gray silty metashale and metaconglomerate; metagartzite clasts are common in the metaconglomerate; as much as 7,315 m thick.
YXra	Red Creek Quartzite (Middle Proterozoic to Upper Archean) - Contains three main rock types: metagartzite, mica schist, and amphibolite; other minor rock types include metadiorite and metacarbonate to marble. Map unit as much as 6,096 m thick.
XWrq	Amphibolite (YXra) - Dark-gray to black, fine- to medium-grained amphibolite composed of strongly foliated to non-foliated metamorphosed mafic rocks, mostly hornblende, intruded into and intimately associated with the Red Creek Quartzite as numerous small bodies in the northeast part of the quadrangle.
XWrm	Metagartzite (XWrq) - Resistant white, gray, tan, and light-green metagartzite.
XWre	Mica schist (XWrm) - Quartz-muscovite schist that grades between metagartzite and mica schist and contains garnet and staurolite.
XWrc	Metadiorite (XWre) - Metamorphosed diorite; epidiorite of previous mappers. Carbonate rock (XWrc) - Metamorphosed carbonate rock along Goslin fault.
Wo	Owiyukuts Complex (Late Archean) - High-grade, metamorphosed potassium-rich granitic gneiss and lesser quartzofeldspathic gneiss; Rb/Sr 2,700 Ma; unknown thickness.

